

# A theory of $\pi/2$ superconducting Josephson junctions

A.Zyuzin

*Max-Planck-Institut für Physik Komplexer Systeme, 01187 Dresden Germany*

*A.F.Ioffe Physical- Technical Institute, 194021, St.Petersburg, Russia*

B.Spivak

*Physics Department University of Washington, Seattle, WA 98195, USA*

## Abstract

We consider theoretically a Josephson junction with a superconducting critical current density which has a random sign along the junction's surface. We show that the ground state of the junction corresponds to the phase difference equal to  $\frac{\pi}{2}$ . Such a situation can take place in superconductor-ferromagnet-superconductor junctions.

Suggested PACS index category: 05.20-y, 82.20-w

The superfluid density connects supercurrent with superfluid velocity in superconductors. In the framework of BCS theory it is a positive quantity. Thus the ground state of a superconductor corresponds to zero superfluid velocity (See for example <sup>[1]</sup>). The question whether and under what circumstances the superfluid density can be negative has been discussed in many theoretical papers. a). It has been shown <sup>[2–5]</sup> that if the distribution function of quasiparticles in superconductors is non-equilibrium the superfluid density can be negative. Negative Josephson coupling has been observed experimentally in nonequilibrium superconductor-normal metal-superconductor junction <sup>[6]</sup>. b). In the presence of magnetic impurities or resonance states with strong electron-electron correlation inside the insulator a superconductor-insulator-superconductor junction could have negative Josephson coupling <sup>[7–10]</sup>. c). In case of disordered superconductor-ferromagnet-superconductor junctions the Josephson coupling averaged over samples has oscillating sign and decays exponentially with the ferromagnetic layer thickness  $L$  <sup>[12]</sup>. Mesoscopic fluctuations of the coupling, however, do not decay exponentially with  $L$ . Therefore at large  $L$  the junctions have couplings with random sign <sup>[13]</sup>. d). At last, a d-superconductor-normal metal-d-superconductor junction can have arbitrary sign depending on the relative orientation of the (superconducting) order parameters in the superconductors <sup>[14,15]</sup>. The fact that the superfluid density is negative indicates that the state with zero phase gradient is unstable. The ground state of a Josephson junction with negative critical current corresponds to a phase difference on the junction which equals  $\pi$ .

In real situation, however, the sign of the superfluid density can fluctuate from point to point inside the sample. In this paper we consider a Josephson junction, with a coupling  $J(\boldsymbol{\rho})$  fluctuating from point to point along the junction's surface and with a random sign. Here  $\boldsymbol{\rho}$  is a coordinate along the junction's surface. We show that in this case the ground state of the system corresponds to a phase difference of the order parameter of  $\frac{\pi}{2}$ .

The total energy of the system can be written as

$$E = E_J + \frac{mn_s}{2} \int d^3\mathbf{r} \left( \mathbf{v}_s(\mathbf{r}) \right)^2 \quad (1)$$

Here  $m$  is the electron mass,  $\mathbf{v}_s(\mathbf{r}) = \frac{1}{m}\nabla\phi(\mathbf{r})$  and  $\phi(\mathbf{r})$  are the superconducting velocity and the phase of the superconducting order parameter in the bulk of the superconductors, ( $\hbar \equiv 1$ ),  $n_s$  is the density of superconducting electrons, which is assumed to be the same in both superconductors,  $E_J$  is the Josephson energy of the junction which can be written as

$$E_J = - \int d^2\boldsymbol{\rho} J(\boldsymbol{\rho}) \cos(\chi(\boldsymbol{\rho})), \quad (2)$$

$\chi(\boldsymbol{\rho}) = \phi(z = +0, \boldsymbol{\rho}) - \phi(z = -0, \boldsymbol{\rho})$  is the jump of the phase of superconducting order parameter at the junction's surface  $z = 0$ . Here  $z$  is the coordinate perpendicular to the junction's surface. In Eqs.1,2 we neglected the thickness of the junction. The second term in Eq.1 corresponds to the energy associated with supercurrents in the bulk of the superconductors. We neglect the magnetic field energy in the bulk of the superconductors assuming that  $J(\boldsymbol{\rho})$  is small and the Josephson penetration length of the magnetic field along the junction's boundary is large.

In the case, when  $J(\boldsymbol{\rho})$  has a random sign, the ground state of the system corresponds to nonuniform distributions of  $\mathbf{v}_s(\mathbf{r}) \neq 0$  and  $\chi(\boldsymbol{\rho}) \neq 0$ . The current conservation law in the bulk of superconductors and at the junction's boundary gives

$$\Delta\phi(\mathbf{r}) = 0 \quad (3)$$

$$\frac{d\phi(\boldsymbol{\rho}, z = +0)}{dz} = \frac{d\phi(\boldsymbol{\rho}, z = -0)}{dz} = \frac{m}{n_s} J(\boldsymbol{\rho}) \sin(\chi(\boldsymbol{\rho})) \quad (4)$$

respectively. Solutions of Eqs.3,4 determine the extrema of the total energy Eqs.1 and 2. At given  $\langle\chi(\boldsymbol{\rho})\rangle$  the minimum of Eq.1 corresponds to some distribution of  $\delta\chi(\boldsymbol{\rho})$ , which is a random sample specific quantity. If  $J(\boldsymbol{\rho})$  is small enough, then  $\delta\chi(\boldsymbol{\rho}) = \chi(\boldsymbol{\rho}) - \langle\chi(\boldsymbol{\rho})\rangle \ll 1$ . Here the bracket  $\langle\rangle$  stands for averaging over the junction's surface.

Substituting  $\chi(\boldsymbol{\rho}) \sim \langle\chi(\boldsymbol{\rho})\rangle$  in left hand side of Eq.4 and solving Eqs.3-4 we obtain an expression for the minimum of the total junction's energy per unite area,

$$E_{eff} = -\langle J(\boldsymbol{\rho}) \rangle \cos\langle\chi\rangle - J_{eff} \sin^2\langle\chi\rangle \quad (5)$$

as a function of  $\langle\chi(\boldsymbol{\rho})\rangle$ , where

$$J_{eff} = \frac{m}{S n_s} \int d\boldsymbol{\rho}_1 d\boldsymbol{\rho}_2 \frac{\langle \delta I(\boldsymbol{\rho}_1) \delta I(\boldsymbol{\rho}_2) \rangle}{2\pi |\boldsymbol{\rho}_1 - \boldsymbol{\rho}_2|} \quad (6)$$

Here  $S$  is the area of the junction. For  $0 < 2J_{eff} < \langle J \rangle$  the junction's energy  $E_{eff}$  has a minimum for  $\langle \chi \rangle = 0$ , while at  $2J_{eff} > \langle J \rangle$ , the minimum corresponds to a nonzero phase difference

$$\langle \chi \rangle^{min} = \arccos\left(\frac{\langle J \rangle}{2J_{eff}}\right), \quad (7)$$

which in the case of  $J_{eff} \gg \langle J \rangle$  gives  $\langle \chi \rangle^{min} = \frac{\pi}{2}$ . The derivation presented above is similar to the case of a ferromagnet-normal metal-ferromagnet junction with a random in sign exchange energy between the ferromagnets <sup>[16]</sup>

Below we consider in more details the case of a superconductor- disordered ferromagnetic metal-superconductor junction shown in Fig.1. We will describe the system by the Hamiltonian

$$H = H_{BCS} + H_T + H_F. \quad (8)$$

where  $H_{BCS}$  is the BCS Hamiltonian of superconducting leads,

$$H_T = t \sum_{i=1,2;\alpha} \int_{S_i} \prod_i d\mathbf{r}_i \left( \Psi_i^+(\mathbf{r}_i, \alpha) \Psi_F(\mathbf{r}, \alpha) + h.c. \right). \quad (9)$$

is the Hamiltonian describing the tunneling between the two superconductors labeled by indexes 1,2, and the ferromagnet labeled by the index  $F$ ; and  $\alpha$  is a spin index. Furthermore,

$$H_F = H_0 + \mathbf{H}\boldsymbol{\sigma} \quad (10)$$

is the Hamiltonian of the ferromagnet. The integration in Eq.9 is taken over the superconductor-ferromagnet surfaces. The effective magnetic field  $\mathbf{H}$  models the ferromagnetic exchange spin splitting,  $\boldsymbol{\sigma}$  are Pauli matrices;  $H_0$  is the hamiltonian of free electron gas in a random potential  $U(\mathbf{r})$ . We assume the following correlation properties of the random potential :  $\langle U(\mathbf{r}) \rangle = 0$  and  $\langle U(\mathbf{r})U(\mathbf{r}') \rangle = \frac{\delta(\mathbf{r}-\mathbf{r}')}{2\pi\nu_0\tau}$ . Here  $\tau$  is the electron elastic mean free time,  $\nu_0$  is electron density of states at the Fermi level.

Let us introduce a nonlocal Josephson coupling density  $J(\boldsymbol{\rho}, \boldsymbol{\rho}')$  as

$$E_J = \int_{S_i} d\boldsymbol{\rho} d\boldsymbol{\rho}' J(\boldsymbol{\rho}, \boldsymbol{\rho}') \cos(\phi(\boldsymbol{\rho}) - \phi(\boldsymbol{\rho}')) \quad (11)$$

where  $\boldsymbol{\rho}$  and  $\boldsymbol{\rho}'$  are coordinates on the surfaces between the ferromagnet and the two superconductors. The integration in Eq.11 is taken over the surfaces  $S$  and  $S'$  of the superconductors.

To lowest order in  $t$  we have

$$J(\boldsymbol{\rho}, \boldsymbol{\rho}') = \frac{T}{2} \sum_{\epsilon_n, \alpha} t^4 \int_{S, S'} d\mathbf{r} d\mathbf{r}' \left[ F_{\alpha, -\alpha}^+(\epsilon_n; \boldsymbol{\rho}; \mathbf{r}) G_{\alpha, \alpha}(\epsilon_n; \mathbf{r}, \boldsymbol{\rho}') F_{\alpha, -\alpha}(\epsilon_n; \boldsymbol{\rho}'; \mathbf{r}') G_{-\alpha, -\alpha}(-\epsilon_n; \mathbf{r}', \boldsymbol{\rho}) + h.c. \right] \quad (12)$$

Here  $F_{\alpha, -\alpha}^+(\epsilon_n; \mathbf{r}_1 \mathbf{r}_1')$  and  $F_{\alpha, -\alpha}(\epsilon_n; \mathbf{r}_1 \mathbf{r}_1')$  are anomalous Green functions of the superconducting leads.  $G_{\alpha, \alpha}(\epsilon_n; \mathbf{r}_1, \mathbf{r}_2)$  are Green functions of electron in ferromagnets;  $\epsilon_n = (2n + 1)\pi T$  is the fermionic Matsubara frequency. A derivation identical to Eq.6 gives

$$J_{eff} = \frac{m}{S n_s} \int d\boldsymbol{\rho} d\boldsymbol{\rho}_1 d\boldsymbol{\rho}' d\boldsymbol{\rho}'_1 \frac{\langle \delta J(\boldsymbol{\rho}, \boldsymbol{\rho}_1) \delta J(\boldsymbol{\rho}', \boldsymbol{\rho}'_1) \rangle}{2\pi |\boldsymbol{\rho} - \boldsymbol{\rho}'|} \sim \frac{\xi m}{\pi n_s} \left( \frac{g}{8\pi\nu_0 D} \right)^4 \left( \frac{D}{2\pi^2 L^2} \right)^2 \ln\left(\frac{\xi}{l}\right). \quad (13)$$

Here  $L$  is the ferromagnet's thickness,  $\xi = \sqrt{D/\Delta}$  is the superconducting coherence length and  $\Delta$  is the modulus of the superconducting order parameter,  $g$  is the dimensionless conductance of superconductor-ferromagnet boundary per unit area, and  $D$  is the electron diffusion constant, which is assumed to be the same in the superconductors and in the ferromagnet. Diagrams which contribute to Eq.13 are shown in Fig.1. We use the conventional diagram technique for the averaging over scattering potential configurations<sup>[17]</sup> and retain the symbol  $\langle \rangle$  to indicate the averaging over configurations of the scattering potential.

In the case of no electron reflection from the superconductor-ferromagnet boundary we can estimate  $J_{eff}$  by taking into account that the mesoscopic fluctuations of the critical current of a SNS junction of the area  $L^2$  is of order of  $\frac{D}{L^2}$ <sup>[11]</sup>. Fluctuations of the Josephson coupling associated with different areas of the junction's surface separated by distances larger than  $L$  are uncorrelated. As a result, we estimate

$$J_{eff} \sim \frac{m}{n_s L} \left( \frac{D}{L^2} \right)^2 L^{-2} \quad (14)$$

It follows from Eq.12 that  $J_{eff}$  decays with  $L$  only as a power law. On the other hand,  $\langle J(\boldsymbol{\rho}) \rangle$  decreases exponentially at  $L > \sqrt{D/H}$  [12] and we have  $|\langle J(\boldsymbol{\rho}) \rangle| \ll J_{eff}$ .

In conclusion we would like to mention that the state discussed above of the superconductor-ferromagnet-superconductor junction is similar to Fulde-Ferrell-Larkin-Ovchinnikov state of bulk superconductors [18,19], when in the presence of the spin splitting energy the superconducting ground state corresponds to a nonuniform distribution of the order parameter. After averaging over the configurations of the impurity potential the order parameter in such a state is significantly suppressed by static disorder. Mesoscopic fluctuations of the order parameter in such a state can however survive the disorder.

Authors kindly acknowledge discussions with P. Fulde.

## REFERENCES

- [1] P.G. de Gennes, Superconductivity of Metals and Alloys”, Addison-Wesley, 1989.
- [2] V.Baru, A.Sukhanov, Pisma Zh.Eksp.Teor.Fiz. **21**, 209, 1975.
- [3] A. Aronov, B. Spivak, Fiz. Tverd. Tela **17**, 2806 (1975), [ Sov. Phys. Solid State **17**, 1874 (1975)]; Fiz.Nizk.Temp.**4** ,1365,1978 [Sov.J.Low Temp.Phys. **4**,641, 1978].
- [4] E.Ginzburg, B.Spivak, Zh. Eksp.Theor.Fiz. **80**, 2013, 1981 [Sov.Phys.JETP. **53**, 1047, 1981].
- [5] B.J. van Wees, K.L.H. Lenssen, C.J.P.M. Harmans, Phys.Rev.B. **44**, 470, 1991.
- [6] J.J.A.Baselmans, et all, Nature **397**(8714), 43, (1999).
- [7] H.Soda,T.Shiba,Prog.Theor.Phys. **41**,25,1969.
- [8] I.O.Kulik Sov.Phys.JETP **22**, 841, 1966.
- [9] L. Bulaevskii, V. Kuzii, A. Sobyenin, Pisma Zh. Eksp. Teor. Fiz. **25**, 314(1977) [JETP Lett. **25**, 7(1977)].
- [10] B.Spivak, S.Kivelson, Phys.Rev.B, **43**, 3740, 1991; S.Kivelson, B.Spivak, Phys.Rev.B, **45**, 10490, 1992.
- [11] B. Altshuler and B. Spivak, Sov.Phys.JETP **65** , 343 1987.
- [12] A.I. Buzdin and M.Yu. Kupriyanov, Sov.Phys. JETP Lett., **53**, 321, 1991.
- [13] F.Zhou, B.Spivak, Phys.Rev.Lett. **80**,5647,1998.
- [14] V.B. Geshkenbein, A.I. Larkin and A. Barone, Phys.Rev.B. **36**, 235, 1987.
- [15] M. Sigrist, T.M. Rice, Rev. Mod. Phys., **67**, 503 1995. Lett., **53**, 321, 1991.
- [16] J.C. Slonczewski, Phys.Rev.Lett. **67**, 3172, 1991.
- [17] A.Abrikosov L.Gorkov D.Dzialoshinskii, ”Methods of quantum field theory in statistical

physics” ,NY, 1963

[18] P. Fulde, R. Ferrell, Phys.Rev.A, **135**, 550. 1964.

[19] A. Larkin, Yu. Ovchinnikov, Sov.Phys. JETP **20**, 762, 1965. Zh. Eksp.Teor.Fiz **47**, 1136, 1964.



# FIGURES




FIG. 1. A schematic picture of the superconductor (S)-ferromagnet (F)- superconductor (S) junction and the diagram for calculation of Eq.13. Thin solid lines inside the ferromagnet correspond to electronic Green functions  $G(\boldsymbol{\rho}, \boldsymbol{\rho}_1)$ , dashed lines correspond to the elastic scattering, thin solid lines inside the superconductors correspond to the anomalous Green function  $F(\boldsymbol{\rho}, \boldsymbol{\rho}')$ .